

Datasheet: 0300-0135

| Description: | GOAT F(ab) ANTI RABBIT IgG (H/L) | | |
|---------------|----------------------------------|--|--|
| Specificity: | IgG (H/L) | | |
| Format: | Purified | | |
| Product Type: | Polyclonal Antibody | | |
| Isotype: | Polyclonal IgG | | |
| Quantity: | 1 mg | | |

Product Details

| Applications | This product has been reported to work in the following applications. This information is derived | | | | | |
|--------------|---|--|--|--|--|--|
| | from testing within our laboratories, peer-reviewed publications or personal communications from | | | | | |
| | the originators. Please refer to references indicated for further information. For general protocol | | | | | |
| | recommendations, please visit www.bio-rad-antibodies.com/protocols. | | | | | |

YesNoNot DeterminedSuggested DilutionELISA•Where this product has not been tested for use in a particular technique this does not necessarily
exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is
recommended that the user titrates the product for use in their own system using the appropriate
negative/positive controls.

| Rabbit | | | |
|--|--|--|--|
| F(ab) fragment of IgG - liquid | | | |
| Purified F(ab) prepared from monospecific affinity-purified IgG by papain digestion followed by chromatographic separation. | | | |
| Phosphate buffered saline | | | |
| 0.01% Sodium Azide (NaN ₃) | | | |
| F(ab) concentration 1.0mg/ml | | | |
| Rabbit IgG whole molecule. | | | |
| UniProt:P01840Related reagentsP01870Related reagentsP01839Related reagentsP01841Related reagentsP01838Related reagentsP01847Related reagents | | | |
| | | | |

| | Entrez Gene: | | | | | |
|-------------------|---|---------------------|--------------------------------|--|--|--|
| | <u>100009097</u> | LOC100009097 | Related reagents | | | |
| | <u>100422811</u> | IGKC2 | Related reagents | | | |
| Specificity | Goat F(ab') anti Rabbit IgG (H+L) antibody recognizes the heavy and light chains of rabbit IgG. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-goat serum. No reaction was observed against anti-papain or anti-goat IgG Fc. | | | | | |
| Storage | Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. | | | | | |
| Shelf Life | 18 months from date of despatch. | | | | | |
| Health And Safety | Material Safety | / Datasheet docur | nentation available at: | | | |
| Information | Material Safety | / Datasheet Docur | mentation #10040 available at: | | | |
| | https://www.bio | o-rad-antibodies.co | pm/uploads/MSDS/10040.pdf | | | |
| Regulatory | For research p | urposes only | | | | |

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (H/L) (602001...) <u>Alk. Phos.</u>, <u>FITC</u> Donkey Anti Goat IgG (H/L) (642001...)<u>Alk. Phos.</u>, <u>Biotin</u>, <u>FITC</u>, <u>HRP</u>, <u>RPE</u>

| North & South | Tel: +1 800 265 7376 | Worldwide | Tel: +44 (0)1865 852 700 | Europe | Tel: +49 (0) 89 8090 95 21 |
|---------------|--------------------------------|-----------|-------------------------------|---------|--------------------------------------|
| America | Fax: +1 919 878 3751 | | Fax: +44 (0)1865 852 739 | | Fax: +49 (0) 89 8090 95 50 |
| | Email: antibody_sales_us@bio-r | rad.com | Email: antibody_sales_uk@bio- | rad.com | Email: antibody_sales_de@bio-rad.com |

'M314379:180412'

Printed on 13 Apr 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint