

Datasheet: 103001

| Description: | GOAT ANTI MOUSE IgG | | |
|---------------|---------------------|--|--|
| Specificity: | IgG | | |
| Format: | Purified | | |
| Product Type: | Polyclonal Antibody | | |
| Isotype: | Polyclonal IgG | | |
| Quantity: | 1 mg | | |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|--------------------|
| Flow Cytometry | | | • | |
| Immunohistology - Frozen | | | | |
| Immunohistology - Paraffin | | | • | |
| ELISA | | | | 1/2000 - 1/4000 |
| Immunoprecipitation | | | • | |
| Western Blotting | | | • | |
| Immunoblotting | | | | |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

| Target Species | Mouse |
|----------------|--|
| Product Form | lg fraction - liquid |
| Preparation | Purified Ig prepared by affinity chromatography on pooled mouse IgG covalently linked to agarose |

Antiserum Preparation Antisera to mouse IgG were raised by repeated immunisations of goats with highly purified antigen.

| Buffer Solution | Borate buffered saline |
|-----------------------------------|----------------------------|
| Preservative Stabilisers | None Present |
| Approx. Protein Concentrations | Ig concentration 1.0 mg/ml |
| External Database | |

External Database Links

UniProt:

P01867 Related reagents
P01865 Related reagents

| | P01863 Related reagents | | | | |
|----------------------------------|---|--------------------|--|------------------------------|--|
| | P01864 | Related | | | |
| | P01868 | Related | | | |
| | P01869 | Related | | | |
| | P03987 | Related | <u>l reagents</u> | | |
| | Entrez Ge | ne: | | | |
| | <u>16016</u> | lghg2b | Related reagents | | |
| | <u>380793</u> | lgh-1a | Related reagents | | |
| | <u>16017</u> | lghg1 | Related reagents | | |
| | <u>16017</u> | lghg1 | Related reagents | | |
| | <u>380793</u> | lgh-1a | Related reagents | | |
| | <u>380793</u> | lgh-1a | Related reagents | | |
| | <u>380795</u> | Al324046 | Related reagents | | |
| Synonyms | lgh-4 | | | | |
| Specificity | | _ | antibody recognizes mouse IgG, recognising d IgG3 as demonstrated by ELISA. | the heavy chain of mouse | |
| | | _ | ntibody has been cross-adsorbed against mo e potential cross-reactivity. | use IgM, mouse IgA and | |
| References | Joimel, U. et al. (2010) Stimulation of angiogenesis resulting from cooperation between macrophages and MDA-MB-231 breast cancer cells: proposed molecular mechanism and effect of tetrathiomolybdate. <u>BMC Cancer. 10: 375.</u> Childs K et al. (2012) Paramyxovirus V proteins interact with the RNA Helicase LGP2 to inhibit RIG-I-dependent interferon induction. <u>J Virol. 86 (7): 3411-21.</u> Moalli, F. et al. (2015) Intravital and whole-organ imaging reveals capture of melanoma-derived antigen by lymph node subcapsular macrophages leading to widespread deposition on follicular | | | | |
| | dendritic cells. Front Immunol. 6: 114. 4. Ramos-Sevillano, E. et al. (2016) PSGL-1 on Leukocytes is a Critical Component of the Host | | | | |
| | Immune Re | esponse aga | ainst Invasive Pneumococcal Disease. <u>PLoS F</u> | Pathog. 12 (3): e1005500. | |
| | 5. Abbate, F. <i>et al.</i> (2016) Acid-sensing ion channel immunoreactivities in the cephalic neuromasts of adult zebrafish. Ann Anat. 207: 27-31. | | | | |
| | or adult zet | oransn. <u>Ann</u> | Affat. 207: 27-31. | | |
| Storage | Store at -20 | 0°C only. | | | |
| | Storage in | frost-free fre | eezers is not recommended. | | |
| | This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature | | | | |
| | the antibod use. | y. Should th | is product contain a precipitate we recommen | d microcentrifugation before | |
| Shelf Life | Please see | label for ex | piry date | | |
| Health And Safety Information | | - | eet documentation #10123 available at: -rad-antibodies.com/uploads/MSDS/10123.pdf | | |
| Regulatory | For researc | ch purposes | only | | |

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) FITC, HRP

Email: antibody_sales_us@bio-rad.com

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

'M316889:180612'

Printed on 20 Jun 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint