

Datasheet: 1065002S

Description:	MOUSE ANTI HUMAN ZAP 70:FITC
Specificity:	ZAP 70
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	SBZAP
Isotype:	lgG1
Quantity:	25 μg

## **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="https://www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm<sup>™</sup> (Product Code <u>BUF09</u>) for this purpose.

Target Species	Human				
Product Form	Purified IgG conj	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FIT			
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)		
	FITC	490	525		
Buffer Solution	Phosphate buffered saline				
Preservative Stabilisers	0.1% Sodium Azide (NaN <sub>3</sub> )				
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml				
Immunogen	Synthetic peptide corresponding to amino acids 280-309 of human ZAP70 conjugated to keyhole limpet hemocyanin (KLH).				
External Database	UniProt:				
Links		Related reagents			

#### **Entrez Gene:**

7535 ZAP70 Related reagents

Synonyms	SRK				
Specificity	Mouse anti Human ZAP 70 antibody, clone SBZAP recognizes human ZAP-70, a 70kDa tyrosine protein kinase which associates with the T-cell receptor (TCR) zeta chain and undergoes phosphorylation following TCR stimulation. ZAP-70 is primarily expressed in T lymphocytes and natural killer cells, where it plays a key role in T-cell receptor (TCR) signalling. Recent studies suggest that ZAP-70 is also expressed in a population of normal immature B-cells and in B-cells from a sub-set of patients with chronic lymphocytic leukaemia.				
References	<ol> <li>Roifman, C.M. <i>et al.</i> (1989) Depletion of CD8+ cells in human thymic medulla results in selective immune deficiency. <u>J Exp Med. 170: 2177 - 2182.</u></li> <li>Arpaia, E. <i>et al.</i> (1994) Defective T cell receptor signaling and CD8+ thymic selection in humans lacking zap-70 kinase. <u>Cell. 76: 947 - 958.</u></li> </ol>				
Storage	Store at +4°C or at -20°C if preferred.  Storage in frost-free freezers is not recommended.  Avoid repeated freezing and thawing as this may denature the antibody. This product should be stored undiluted.  This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.				
Shelf Life	Please see label for expiry date.				
Health And Safety Information	Material Safety Datasheet documentation #10303 available at:  10303: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10303.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10303.pdf</a>				
Regulatory	For research purposes only				

# **Related Products**

## **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

## **Recommended Useful Reagents**

LEUCOPERM™ (BUF09)
LEUCOPERM™ (BUF09B)
LEUCOPERM™ (BUF09C)
HUMAN SEROBLOCK (BUF070A)

HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_us@bio-rad.com

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

'M310480:170926'

Printed on 11 Jul 2018