

Datasheet: 2150-1470

Description:	RABBIT ANTI MOUSE COLLAGEN IV
Specificity:	COLLAGEN IV
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	■			1/500
Immunohistology - Paraffin	■			1/500
ELISA	■			1/2000
Immunoprecipitation			■	
Western Blotting			■	
Immunofluorescence	■			1/40

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species

Mouse

Species Cross Reactivity

Reacts with: Orangutan, Rat

N.B. Antibody reactivity and working conditions may vary between species.

Product Form

Purified Ig - liquid

Preparation

Purified IgG prepared by antigen column chromatography

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers

0.09% Sodium Azide (NaN₃)

Immunogen

Collagen IV purified from mouse EHS tumor.

External Database Links

UniProt:

- | | |
|--------|----------------------------------|
| P02463 | Related reagents |
| P08122 | Related reagents |
| Q9QZS0 | Related reagents |

Q9QZR9	Related reagents
Q80V57	Related reagents
Q6PFB1	Related reagents

Entrez Gene:

12826	Col4a1	Related reagents
12827	Col4a2	Related reagents
12828	Col4a3	Related reagents
12829	Col4a4	Related reagents

Specificity

Rabbit anti Mouse Collagen IV antibody recognizes mouse collagen type IV. Collagen IV is a 1682 amino acid ~160 kDa (predicted) matrix protein and major component of glomerular basement membranes. Multiple isoforms exist each capable of forming triple helical structures with two other chains to form the type IV collagen network. The collagen IV alpha chain can be cleaved between residues 1444-1445 to yield the c-terminal 225 amino acid, ~28 kDa arresten fragment, collagen α2(IV) yields a c-terminal canstatin fragment while Collagen α3(IV) yeilds a tumstatin fragment. Collagen IV bears a single [collagen IV NC1](#) (C-terminal non-collagenous) domain ([UniProt: Q9QZR9](#)).

Mutations in collagen IV genes have been implicated in inherited nephropathies and potentially in cystic kidney disease and intracranial aneurysms ([Plaisier et al. 2007](#)).

Rabbit anti Mouse Collagen IV antibody has been successfully employed for the detection of collagen IV by immunofluorescence and immunohistochemistry in mice ([Tang et al. 2010](#)), rats ([Shen et al. 2014](#)) and orangutan ([Bredies et al. 2013](#)).

The following cross reactivities have been observed:

Mouse type IV	100%
Mouse types I, II & III	<0.1%
Human types IV & V	<0.1%
Mouse fibronectin & laminin	<0.1%

References

1. Tang, Z. *et al.* (2010) Survival effect of PDGF-CC rescues neurons from apoptosis in both brain and retina by regulating GSK3beta phosphorylation. [J Exp Med. 207: 867-80.](#)
2. Fantin, A. *et al.* (2010) Tissue macrophages act as cellular chaperones for vascular anastomosis downstream of VEGF-mediated endothelial tip cell induction. [Blood.116: 829-40.](#)
3. Shen, W. *et al.* (2014) Systemic Administration of Erythropoietin Inhibits Retinopathy in RCS Rats. [PLoS One. 9: e104759.](#)
4. Bredies, K. *et al.* (2013) Computer-assisted counting of retinal cells by automatic segmentation after TV denoising. [BMC Ophthalmol. 13: 59.](#)
5. Rubin, A.N. *et al.* (2010) The germinal zones of the basal ganglia but not the septum generate GABAergic interneurons for the cortex. [J Neurosci. 30: 12050-62.](#)
6. Weinl, C. *et al.* (2014) Elk3 deficiency causes transient impairment in post-natal retinal vascular development and formation of tortuous arteries in adult murine retinae. [PLoS One. 9: e107048.](#)
7. Stenzel, D. *et al.* (2011) Integrin-dependent and -independent functions of astrocytic fibronectin in retinal angiogenesis. [Development. 138: 4451-63.](#)
8. Zuercher, J. *et al.* (2012) Norrin stimulates cell proliferation in the superficial retinal vascular plexus and is pivotal for the recruitment of mural cells. [Hum Mol Genet. 21: 2619-30.](#)
9. Arnold, T.D. *et al.* (2012) Defective retinal vascular endothelial cell development as a consequence of impaired integrin αVβ8-mediated activation of transforming growth factor-β. [J](#)

Neurosci. 32: 1197-206.

10. Chen, M. *et al.* (2010) Immune activation in retinal aging: a gene expression study. Invest Ophthalmol Vis Sci. 51: 5888-96.
11. Kojima, T. *et al.* (2007) Proangiogenic role of ephrinB1/EphB1 in basic fibroblast growth factor-induced corneal angiogenesis. Am J Pathol. 170: 764-73.
12. Ma, S. *et al.* (2012) Ric-8a, a Guanine Nucleotide Exchange Factor for Heterotrimeric G Proteins, Regulates Bergmann Glia-Basement Membrane Adhesion during Cerebellar Foliation. J Neurosci. 32: 14979-93.
13. Dulauroy, S. *et al.* (2012) Lineage tracing and genetic ablation of ADAM12(+) perivascular cells identify a major source of profibrotic cells during acute tissue injury. Nat Med. 18: 1262-70.
14. Edgar, K. *et al.* (2012) eNOS Overexpression Exacerbates Vascular Closure in the Obliterative Phase of OIR and Increases Angiogenic Drive in the Subsequent Proliferative Stage. Invest Ophthalmol Vis Sci. 53: 6833-50.
15. Lange, C.A. *et al.* (2012) Von Hippel-Lindau protein in the RPE is essential for normal ocular growth and vascular development. Development. 139: 2340-50.
16. Li, W. and Mukouyama, Y.S. (2011) Whole-mount immunohistochemical analysis for embryonic limb skin vasculature: a model system to study vascular branching morphogenesis in embryo. J Vis Exp. 51: pii: 2620.
17. Lutter, S. *et al.* (2012) Smooth muscle-endothelial cell communication activates Reelin signaling and regulates lymphatic vessel formation. J Cell Biol. 197: 837-49.
18. McKenzie, J.A. *et al.* (2012) Apelin is required for non-neovascular remodeling in the retina. Am J Pathol. 180: 399-409.
19. Powner, M.B. *et al.* (2012) Visualization of gene expression in whole mouse retina by in situ hybridization. Nat Protoc. 7: 1086-96.
20. Schulz, C. *et al.* (2012) A lineage of myeloid cells independent of Myb and hematopoietic stem cells. Science. 336: 86-90.
21. Scott, A. *et al.* (2010) Astrocyte-derived vascular endothelial growth factor stabilizes vessels in the developing retinal vasculature. PLoS One. 5: e11863.
22. Takagi, N. *et al.* (2012) Mineralocorticoid Receptor Blocker Protects against Podocyte-Dependent Glomerulosclerosis. Nephron Extra. 2: 17-26.
23. Chen, M. *et al.* (2013) Age- and light-dependent development of localised retinal atrophy in CCL2(-/-)CX3CR1(GFP/GFP) mice. PLoS One. 8: e61381.
24. Luhmann UF. *et al.* (2005) Berger W. Role of the Norrie disease pseudoglioma gene in sprouting angiogenesis during development of the retinal vasculature. Invest Ophthalmol Vis Sci. 46: 3372-82.
25. Scott, A. *et al.* (2014) Quantification of vascular tortuosity as an early outcome measure in oxygen induced retinopathy (OIR). Exp Eye Res. 120: 55-60.
26. Yukiura H *et al.* (2015) Autotaxin overexpression causes embryonic lethality and vascular defects. PLoS One. 10 (5): e0126734.
27. Williams, J.A. *et al.* (2016) Regulation of C3 Activation by the Alternative Complement Pathway in the Mouse Retina. PLoS One. 11 (8): e0161898.
28. Piñero G *et al.* (2016) Lithium Reversibly Inhibits Schwann Cell Proliferation and Differentiation Without Inducing Myelin Loss. Mol Neurobiol. Dec 5. [Epub ahead of print]
29. Gurnik, S. *et al.* (2016) Angiopoietin-2-induced blood-brain barrier compromise and increased stroke size are rescued by VE-PTP-dependent restoration of Tie2 signaling. Acta Neuropathol. 131 (5): 753-73.
30. Misra, A. *et al.* (2016) Integrin β 3 inhibition is a therapeutic strategy for supravalvular aortic stenosis. J Exp Med. 213 (3): 451-63.
31. Wu, W.K. *et al.* (2015) IL-4 regulates specific Arg-1(+) macrophage sFlt-1-mediated inhibition of angiogenesis. Am J Pathol. 185 (8): 2324-35.
32. Toffoli, B. *et al.* (2017) Nephropathy in Pparg-null mice highlights PPAR γ systemic activities in metabolism and in the immune system. PLoS One. 12 (2): e0171474.
33. Yanagida, K. *et al.* (2017) Size-selective opening of the blood-brain barrier by targeting

endothelial sphingosine 1-phosphate receptor 1. [Proc Natl Acad Sci U S A. 114 \(17\): 4531-6.](#)

34. Fernández-robredo, P. et al. (2017) Neuropilin 1 Involvement in Choroidal and Retinal Neovascularisation. [PLoS One. 12 \(1\): e0169865.](#)

Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...) [FITC](#)

Sheep Anti Rabbit IgG (STAR35...) [RPE](#)

Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)

Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)

Sheep Anti Rabbit IgG (2AB02...) [Biotin](#)

Sheep Anti Rabbit IgG (STAR36...) [DyLight®488](#), [DyLight®549](#), [DyLight®649](#),
[DyLight®680](#), [DyLight®800](#)

Recommended Useful Reagents

[ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025A\)](#)

[ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025C\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M318334:180718'

Printed on 01 Aug 2018