

Datasheet: 2150-1948

Description:	RABBIT ANTI RAT COLLAGEN III
Specificity:	COLLAGEN III
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	50 µg

## **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	<b>Not Determined</b>	Suggested Dilution
Immunohistology - Frozen				1/20
Immunohistology - Paraffin				
ELISA				1/3000
Western Blotting				
Immunofluorescence				

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls.

Target Species	Rat
Product Form	Purified IgG - lyophilised
Reconstitution	Reconstitute with 0.5 ml distilled water.

Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution.

# Antiserum Preparation Antiserum to rat collagen type III was raised by repeated immunisation of rabbits with highly

purified antigen. Purified IgG was prepared from whole serum by affinity chromatography on

immobilized rat collagen type III.

Buffer Solution	Phosphate/Borate buffered saline.
Preservative Stabilisers	1% Mannitol 1% Dextran
Approx. Protein Concentrations	0.1 mg/ml after reconstitution
Immunogen	Collagen type III from rat tail tendon.

**External Database** Links

UniProt:

P13941 Related reagents

**Entrez Gene:** 

84032 Col3a1 Related reagents

Specificity

Rabbit anti Rat Collagen III antibody recognizes rat collagen type III, binding to both native and heat denatured collagen type III. This antibody is reactive with rat kidney, liver, skin and heart, but does not stain basement membranes.

The antiserum was cross-absorbed with immobilized rat serum proteins.

References

1. Jones, H.B. et al. (2013) Quantitative histopathological assessment of retardation of islets of langerhans degeneration in rosiglitazone-dosed obese ZDF rats using combined insulin and collagens (I and III) immunohistochemistry with automated image analysis and statistical modeling. Toxicol Pathol. 41 (3): 425-44.

2. Dankers P. Y.W. et al. (2015) Convenient formulation and application of a supramolecular ureidopyrimidinone modified poly(ethylene glycol) carrier for intrarenal growth factor delivery European Polymer Journal. Jul 2 [Epub ahead of Print]

**Storage** 

Prior to reconstitution store at +4°C. After reconstitution store at -20°C.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody.

**Shelf Life** 12 months from date of reconstitution.

**Health And Safety** Information

Material Safety Datasheet documentation #10325 available at: 10325: https://www.bio-rad-antibodies.com/uploads/MSDS/10325.pdf

Regulatory For research purposes only

### Related Products

### **Recommended Secondary Antibodies**

Sheep Anti Rabbit IgG (STAR34...) **FITC** Sheep Anti Rabbit IgG (STAR35...) **RPE** Goat Anti Rabbit IgG (H/L) (STAR124...) HRP

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

Sheep Anti Rabbit IgG (2AB02...)

Sheep Anti Rabbit IgG (STAR36...) DyLight®488, DyLight®549, DyLight®649,

DyLight®680, DyLight®800

America

North & South Tel: +1 800 265 7376

Fax: +1 919 878 3751 Email: antibody\_sales\_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700 Europe

Fax: +44 (0)1865 852 739

Email: antibody\_sales\_uk@bio-rad.com

Fax: +49 (0) 89 8090 95 50

Tel: +49 (0) 89 8090 95 21

Email: antibody\_sales\_de@bio-rad.com

'M316991:180615'

Printed on 21 Jun 2018