

Datasheet: 4660-1140

| Description: | GOAT ANTI HUMAN GLUCAGON (N-TERMINAL) |
|---------------|---------------------------------------|
| Specificity: | GLUCAGON (N-TERMINAL) |
| Format: | Purified |
| Product Type: | Polyclonal Antibody |
| lsotype: | Polyclonal IgG |
| Quantity: | 1 ml |

Product Details

| Applications | - | | | | lowing applications. This | | | |
|-----------------------------------|--|--------------------------|-----|-----|---------------------------|--------------------|--|--|
| | from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols. | | | | | | | |
| | | | | | | | | |
| | Flow Cytometry | | 105 | 110 | | ouggested blidtion | | |
| | | Immunohistology - Frozen | | | | | | |
| | Immunohistolog | | • | | | | | |
| | ELISA | , | • | | | | | |
| | Western Blotting | 3 | • | | | | | |
| | Western blocking Where this product has not been tested for use in a particular technique this does not necessaril | | | | | | | |
| | exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls. | | | | | | | |
| | | | | | | | | |
| Target Species | Human | | | | | | | |
| Species Cross Reactivity | Reacts with: Pig N.B. Antibody reactivity and working conditions may vary between species. | | | | | | | |
| Product Form | Purified IgG - liquid | | | | | | | |
| Preparation | Purified IgG prepared by affinity chromatography on Protein G | | | | | | | |
| Buffer Solution | Phosphate buffered saline | | | | | | | |
| Preservative Stabilisers | 0.09% Sodium Azide (NaN ₃) | | | | | | | |
| Approx. Protein Concentrations | IgG concentration 5.0 mg/ml | | | | | | | |
| Immunogen | Porcine glucagon, amino acids 1-12, conjugated with bovine thyroglobulin. | | | | | | | |
| External Database | UniProt: | | | | | | | |
| Links | J | | | | | | | |

Entrez Gene: <u>2641</u> GCG **Related reagents** Specificity Goat anti Human glucagon antibody recognizes the pancreatic polypeptide hormone glucagon, whose actions are antagonistic to insulin. Glucagon is produced by A cells within the Islets of Langerhans in the pancreas. Tumors arising from glucagon producing cells are termed glucagonomas, and this antibody may be useful in identifying such tumours. References 1. Li, S. et al. (2010) SEL1L deficiency impairs growth and differentiation of pancreatic epithelial cells. BMC Dev Biol. 10: 19. Storage Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the protein. Should this product contain a precipitate we recommend microcentrifugation before use. Shelf Life 18 months from date of despatch. **Health And Safety** Material Safety Datasheet documentation #10040 available at: Information 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) FITC, HRP

| North & South | Tel: +1 800 265 7376 | Worldwide | Tel: +44 (0)1865 852 700 | Europe | Tel: +49 (0) 89 8090 95 21 |
|---------------|---------------------------------|-----------|---------------------------------|--------|--------------------------------------|
| America | Fax: +1 919 878 3751 | | Fax: +44 (0)1865 852 739 | | Fax: +49 (0) 89 8090 95 50 |
| | Email: antibody_sales_us@bio-ra | d.com | Email: antibody_sales_uk@bio-ra | d.com | Email: antibody_sales_de@bio-rad.com |

'M271475:150528'

Printed on 02 May 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint