

## Datasheet: AAI19F

| Description:  | SHEEP ANTI BOVINE IgM:FITC |  |  |
|---------------|----------------------------|--|--|
| Specificity:  | IgM                        |  |  |
| Format:       | FITC                       |  |  |
| Product Type: | Polyclonal Antibody        |  |  |
| Isotype:      | Polyclonal IgG             |  |  |
| Quantity:     | 1 mg                       |  |  |

## **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

|                            | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|--------------------|
| Flow Cytometry             | -   |    |                | 1/20 - 1/100       |
| Immunohistology - Frozen   | -   |    |                | 1/50 - 1/500       |
| Immunohistology - Paraffin |     |    | •              |                    |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

| Target Species | Bovine   |                         |                   |                        |
|----------------|--|-------------------------|-------------------|------------------------|
| Product Form   | Purified IgG fraction conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liqui |                         |                   | omer 1 (FITC) - liquid |
| Max Ex/Em      | <b>Fluorophore</b><br>FITC   | Excitation Max (nm) 490 | Emission Max (nm) |                        |

**Antiserum Preparation** Antisera to bovine IgM were raised by repeated immunisation of sheep with highly purified antigen. Purified IgG prepared by affinity chromatography.

| <b>Buffer Solution</b>         | Phosphate buffered saline  |
|--------------------------------|--|
| Preservative<br>Stabilisers    | 0.09% Sodium Azide 0.2% Bovine Serum Albumin   |
| Approx. Protein Concentrations | IgG concentration 1.0 mg/ml  |
| Immunogen                      | Purified bovine IgM.   |
| Specificity                    | <b>Sheep anti bovine IgM</b> recognizes bovine IgM and shows no cross-reactivity with other bovine immunoglobulin classes in immunoelectrophoresis. Sheep anti bovine IgM may cross-react with IgM from other species. |

## References

- 1. Nebl, T. *et al.* (2002) Proteomic analysis of a detergent-resistant membrane skeleton from neutrophil plasma membranes. <u>J Biol Chem. 277 (45): 43399-409.</u>
- 2. Assad, A. *et al.* (2012) Immunophenotyping and characterization of BNP colostra revealed pathogenic alloantibodies of IgG1 subclass with specifity to platelets, granulocytes and monocytes of all maturation stages. <u>Vet Immunol Immunopathol. 147: 25-34.</u>
- 3. Hamsten, C. *et al.* (2009) Recombinant surface proteomics as a tool to analyze humoral immune responses in bovines infected by *Mycoplasma mycoides* subsp. *mycoides* small colony type. <u>Mol Cell Proteomics. 8: 2544-54.</u>
- 4. Mansilla FC *et al.* (2015) Safety and immunogenicity of a soluble native *Neospora caninum* tachyzoite-extract vaccine formulated with a soy lecithin/β-glucan adjuvant in pregnant cattle. <u>Vet Immunol Immunopathol. 165 (1-2): 75-80.</u>
- 5. Hossain, M.M. *et al.* (2016) Multiplex Detection of IgG and IgM to Rift Valley Fever Virus Nucleoprotein, Nonstructural Proteins, and Glycoprotein in Ovine and Bovine. <u>Vector Borne Zoonotic Dis. Jul 5. [Epub ahead of print]</u>
- 6. Van Meulder, F. *et al.* (2015) Analysis of the protective immune response following intramuscular vaccination of calves against the intestinal parasite *Cooperia oncophora*. <u>Int J Parasitol. 45 (9-10):</u> 637-46.
- 7. Jankowska, A. *et al.* (2016) Humoral and cellular immune response to *Histophilus somni* recombinant heat shock protein 60 kDa in farm animals <u>Veterinární Medicína. 60 (No. 11): 603-12</u>.

  8. Trotta, M. *et al.* (2015) Simultaneous immunization of cattle with foot-and-mouth disease (FMD) and live anthrax vaccines do not interfere with FMD booster responses <u>Trials in Vaccinology. 4:</u> 38-42.

Storage Store at +4°C. DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Europe

Shelf Life 18 months from date of despatch.

Health And Safety Information Material Safety Datasheet documentation available at:
Material Safety Datasheet Documentation #10041 available at:
<a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf</a>

**Regulatory** For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Fax: +1 919 878 3751

 ${\bf Email: antibody\_sales\_us@bio-rad.com}$ 

Worldwide Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody\_sales\_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody sales de@bio-rad.com

'M314646:180412'

Printed on 13 Apr 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint