

Datasheet: AHP1278

Description:	RABBIT ANTI HUMAN CTGF
Specificity:	CTGF
Other names:	CONNECTIVE TISSUE GROWTH FACTOR
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			•	
Immunohistology - Frozen				
Immunohistology - Paraffin (1)	•			
ELISA	•			0.5 - 2.0ug/ml
Western Blotting	•			0.1 - 0.2ug/ml
Immunofluorescence				
Functional Assays	•			6.0 - 9.0ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species	Human	
Product Form	Purified IgG - lyophilised	
Reconstitution	Reconstitute with 0.1ml distilled water. For long term storage the ad recommended. NB: For functional studies do not add azide.	dition of 0.09% sodium azide is
Antiserum Preparation	Antisera to human CTGF were raised by repeated immunisations of antigen. Purified IgG was prepared from whole serum by affinity chr	3 , .
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	None present	
Approx. Protein Concentrations	IgG concentration 1.0mg/ml after reconstitution.	

Immunogen	Recombinant human CTGF.
External Database	H. P. A
Links	UniProt:
	P29279 Related reagents
	Entrez Gene:
	1490 CTGF Related reagents
Synonyms	CCN2, HCS24, IGFBP8
Specificity	Rabbit anti Human CTGF antibody recognizes human CTGF (connective tissue growth factor), a
	323 amino acid ~38 kDa polypeptide member of the CCN family of secreted cysteine rich
	regulatory proteins. CTGF is produced by vascular endothelial and umbilical vein cells, and
	interacts with several other growth factors, including transforming growth factor beta (TGF-beta),
	vascular endothelial growth factor (VEGF) and bone morphogenetic proteins (BMPs).
	CTGF is a heparin-binding pleiotropic growth factor involved in chondrogenesis and potentially
	skeletogenesis. CTGF promotes the adhesion of epithelial cells and fibroblasts, stimulates integrin
	expression, and also promotes endothelial cell survival, migration and adhesion, implicating CTGF
	in angiogenesis. CTGF induces extracellular matrix formation, including the deposition of excessive
	collagen in fibrotic diseases, and has also been implicated in wound healing.
ELISA	This purified human CTGF antibody may ne used in an indirect ELISA or as the capture reagent in
	a sandwich ELISA with our biotinylated human CTGF antibody (AHP1278B) as the detection
	reagent and recombinant human CTGF (PHP174) as the standard.
References	1. Waddell, J.M. et al. (2011) CTGF expression is up-regulated by PROK1 in early pregnancy and
	influences HTR-8/Svneo cell adhesion and network formation. Hum Reprod. 26 (1): 67-75.
	2. Westermann, D. et al. (2009) Gene deletion of the kinin receptor B1 attenuates cardiac
	inflammation and fibrosis during the development of experimental diabetic cardiomyopathy.
	<u>Diabetes. 58: 1373-81.</u>
	3. Vasilieva, O.V. et al. (2016) Connective tissue growth factor (CTGF) in the human dermis
	through ontogenesis. Russian Journal of Developmental Biology. 47 (2): 63-68.
Further Reading	1. Moussad, E.E. & Brigstock, D.R. (2000) Connective tissue growth factor: what's in a name? Mol
	Genet Metab. 71 (1-2): 276-92.
Storage	Prior to reconstitution store at +4°C.
	After reconstitution store at -20°C.
	Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid
	repeated freezing and thawing as this may denature the antibody. Should this product contain a
	precipitate we recommend microcentrifugation before use.
Shelf Life	12 months from date of reconstitution.
Health And Safety	Material Safety Datasheet documentation #10162 available at:
Information	10162: https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf
momation	

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...) FITC
Sheep Anti Rabbit IgG (STAR35...) RPE
Goat Anti Rabbit IgG (H/L) (STAR124...) HRP

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

Sheep Anti Rabbit IgG (2AB02...) Biotin

Sheep Anti Rabbit IgG (STAR36...) <u>DyLight®488</u>, <u>DyLight®549</u>, <u>DyLight®649</u>,

DyLight®680, DyLight®800

Recommended Useful Reagents

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A)

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025C)

TidyBlot™ WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

'M279278:160115'

Printed on 20 Jun 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint