

## Datasheet: AHP455G

<b>Description:</b>	RABBIT ANTI HUMAN INTERLEUKIN-17
<b>Specificity:</b>	IL-17
<b>Other names:</b>	IL-17A
<b>Format:</b>	Purified
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin (1)	▪			0.25ug/ml
ELISA	▪			0.5 - 2.0ug/ml
Immunoprecipitation			▪	
Western Blotting	▪			0.1 - 0.2ug/ml
Immunofluorescence	▪			
Functional Assays (2)	▪			0.93 - 1.3ug/ml

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) **This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.**

(2) **To yield one-half maximal inhibition [ND50] of the biological activity of hIL-17A (50.0 ng/ml).**

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG - lyophilised
<b>Reconstitution</b>	<p>Reconstitute with 0.1ml distilled water</p> <p>Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution. For long term storage the addition of 0.09% sodium azide is recommended.</p> <p>N.B. For functional studies do not add sodium azide</p>
<b>Antiserum Preparation</b>	Antisera to human IL-17 were raised by repeated immunisation of rabbits with highly purified antigen. Purified IgG prepared by antigen affinity chromatography
<b>Buffer Solution</b>	Phosphate buffered saline

<b>Preservative Stabilisers</b>	None present
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml after reconstitution
<b>Immunogen</b>	<a href="#">Recombinant human IL-17</a>
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">Q16552</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">3605</a>    IL17A    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	CTLA8, IL17
<b>Specificity</b>	<p><b>Rabbit anti Human Interleukin-17 antibody</b> recognizes human IL-17 (Interleukin-17/17A), also known as Cytotoxic T-lymphocyte-associated antigen 8 or CTLA-8. Interleukin-17 is a homodimer of two 132 amino acid chains resulting in a ~32 kDa disulfide-linked cytokine belonging to the IL-17 family (classified as IL-17A to IL-17F) which signals through the IL-17 receptor (IL-17R/CDw217).</p> <p>IL-17 is produced primarily by activated CD4+ T cells and induces a variety of target cells to secrete inflammatory cytokines, including IL-1beta, IL-6, TNFalpha, IFNgamma and granulocyte colony-stimulating factor, as well as synergizing with other inflammatory cytokines and agonists. The role of IL-17 in bone metabolism has become a focus area for studies into Rheumatoid Arthritis, in which levels of IL-17 are significantly increased and may contribute to bone destruction (<a href="#">Ziolkowska et al. 2000</a>).</p>
<b>ELISA</b>	This purified human IL-17 antibody may be used in an ELISA with <a href="#">recombinant human IL-17</a> (PHP098) as the standard.
<b>Western Blotting</b>	This antibody may be used in Western Blotting under either reducing or non-reducing conditions with <a href="#">recombinant human IL-17</a> (PHP098) as the positive control.
<b>References</b>	1. Parsonage, G. <i>et al.</i> (2008) Prolonged, granulocyte-macrophage colony-stimulating factor-dependent, neutrophil survival following rheumatoid synovial fibroblast activation by IL-17 and TNFalpha. <a href="#">Arthritis Res Ther. 10: R47.</a>
<b>Further Reading</b>	1. Gaffen, S.L. (2004) Biology of recently discovered cytokines; Interleukin-17 – a unique inflammatory cytokine with roles in bone biology and arthritis. <a href="#">Arth. Res. and Therapy. 6: 240-247.</a>
<b>Storage</b>	<p>Prior to reconstitution store at +4°C. Following reconstitution store at -20°C.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
<b>Shelf Life</b>	12 months from date of reconstitution.

**Health And Safety Information** Material Safety Datasheet documentation #10162 available at:  
10162: <https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...) [FITC](#)  
Sheep Anti Rabbit IgG (STAR35...) [RPE](#)  
Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)  
Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)  
Sheep Anti Rabbit IgG (2AB02...) [Biotin](#)  
Sheep Anti Rabbit IgG (STAR36...) [DyLight@488](#), [DyLight@549](#), [DyLight@649](#),  
[DyLight@680](#), [DyLight@800](#)

### Recommended Positive Controls

[RECOMBINANT HUMAN INTERLEUKIN-17 \(PHP098\)](#)

### Recommended Useful Reagents

[ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025A\)](#)  
[ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025C\)](#)  
[TidyBlot™ WESTERN BLOT DETECTION REAGENT:HRP \(STAR209P\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

'M318043:180718'

**Printed on 01 Aug 2018**

---

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)