

## Datasheet: AHP933

| Description:  | Description: RABBIT ANTI BIM |  |
|---------------|------------------------------|--|
| Specificity:  | BIM                          |  |
| Other names:  | BOD                          |  |
| Format:       | Purified                     |  |
| Product Type: | Polyclonal Antibody          |  |
| Isotype:      | Polyclonal IgG               |  |
| Quantity:     | 0.1 mg                       |  |
|               |                              |  |

### **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="https://www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

|                                | Yes | No | Not Determined | Suggested Dilution |
|--------------------------------|-----|----|----------------|--------------------|
| Flow Cytometry                 |     |    | •              |                    |
| Immunohistology - Frozen       |     |    | •              |                    |
| Immunohistology - Paraffin (1) |     |    |                |                    |
| ELISA                          |     |    | •              |                    |
| Immunoprecipitation            |     |    | •              |                    |
| Western Blotting               |     |    |                | 1/1000             |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

| Target Species                    | Human  |                            |
|-----------------------------------|--|----------------------------|
| Species Cross<br>Reactivity       | Based on sequence similarity, is expected to react with:Mouse, Rat <b>N.B.</b> Antibody reactivity and working conditions may vary between s | pecies.                    |
| Product Form                      | Purified IgG - liquid  |                            |
| Antiserum Preparation             | Antisera to human Bim were raised by repeated immunisations of ra<br>antigen. Purified IgG prepared by affinity chromatography.              | bbits with highly purified |
| Buffer Solution                   | Phosphate buffered saline  |                            |
| Preservative<br>Stabilisers       | 0.02% Sodium Azide   |                            |
| Approx. Protein<br>Concentrations | IgG concentration 1.0 mg/ml.   |                            |

| Immunogen                        | Synthetic peptide corresponding to amino acids 22 to 40 of human Bim.   |
|----------------------------------|---|
| External Database<br>Links       | UniProt: O43521 Related reagents  Entrez Gene: 10018 BCL2L11 Related reagents   |
| Synonyms                         | BIM   |
| Specificity                      | Rabbit anti Human Bim antibody recognizes human Bcl-2 interacting mediator of cell death (Bim), also known as Bod.  |
|                                  | Bim is a pro-apoptotic protein within the Bcl-2 family, which functions within the intrinsic apotosis pathway. Bim is known to interact with pro-survival proteins including Bcl-2, Bcl-XL and Bcl-w.   |
|                                  | Bim is expressed in a variety of cell types, where it is maintained in an inactive conformation by phosphorylation or through binding to the microtubule-associated dynein motor complex. Upon certain apoptotic stimuli, Bim is released from microtubules and mediates caspase-dependent apoptosis. |
| Western Blotting                 | AHP933 detects a band of approximately 22kDa in K562 and A549 whole cell lysates.   |
| Storage                          | Store at +4°C or at -20°C if preferred.  Storage in frost-free freezers is not recommended.  This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.   |
| Shelf Life                       | 12 months from date of despatch.  |
| Health And Safety<br>Information | Material Safety Datasheet documentation #10040 available at: 10040: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</a>   |
| Regulatory                       | For research purposes only  |

# **Related Products**

# **Recommended Secondary Antibodies**

Sheep Anti Rabbit IgG (STAR34...) FITC
Sheep Anti Rabbit IgG (STAR35...) RPE
Goat Anti Rabbit IgG (H/L) (STAR124...) HRP

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

Sheep Anti Rabbit IgG (2AB02...) Biotin

Sheep Anti Rabbit IgG (STAR36...) <u>DyLight®488</u>, <u>DyLight®549</u>, <u>DyLight®649</u>,

DyLight®680, DyLight®800

### **Recommended Useful Reagents**

TidyBlot™ WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751 Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_de@bio-rad.com

Email: antibody\_sales\_us@bio-rad.com

Email: antibody\_sales\_uk@bio-rad.com

'M285709:160408'

Printed on 30 Apr 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint