

Datasheet: BUF049A

Description:	HISPEC ASSAY DILUENT
Name:	HISPEC ASSAY DILUENT
Format:	Ready To Use
Product Type:	Accessory Reagent
Clone:	N/A
Quantity:	50 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
ELISA	▪			
Immunoassay	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Product Form Ready to use assay diluent buffer

Preservative Stabilisers 0.1% Proclin™ 300

Intended Use HISPEC assay diluent works to reduce cross reactivity, non-specific binding and matrix effects in immunoassays such as ELISA, EIA, Western blotting, immuno-PCR, protein arrays, multianalyte immunoassays and immunohistochemistry.

The effect of HISPEC assay diluent is dependent on assay system and the antibodies used. HISPEC assay diluent is used instead of a sample buffer or antibody dilution buffer within the immunoassay protocol.

HISPEC assay diluent is unsuitable for blocking of surfaces. For blocking of surfaces we recommend one of the blocking solutions within the Bio-Rad range – see “Useful reagents” below.

Examples of use:

- ELISA: use as a dilution buffer for the specimen and for the detection antibodies
- Western blotting: use as dilution buffer for primary and secondary antibodies
- Immunohistochemistry: use as dilution buffer for primary and secondary antibodies
- Protein arrays: use as dilution buffer for specimen and for the detection antibodies

Instructions For Use The buffer should be mixed thoroughly immediately before use.
Dilution of the samples:

The standards and samples for ELISA and protein arrays can be diluted with HISPEC assay diluent at 1:2 or higher. Standards and samples should be treated identically.

Dilution of antibodies:

Antibodies can be diluted with HISPEC assay diluent in a user-defined manner, depending on the recommendation of the antibody supplier.

Expected results:

In some cases a reduction in assay signal intensity may be observed when using HISPEC assay diluent. This does not result in a reduction in assay sensitivity due to the accompanying reduction in non-specific binding.

Signal intensity may be increased in some assay systems by use of higher concentrations of antibodies, or by dilution of the HISPEC assay diluent in distilled water (1:2) prior to use. However, these adjustments may also result in some associated increase in non-specific binding.

References

1. Eriksson, O. *et al.* (2017) Pancreatic imaging using an antibody fragment targeting the zinc transporter type 8: a direct comparison with radio-iodinated Exendin-4. [Acta Diabetol. Oct 24 \[Epub ahead of print\].](#)

Storage

Store at +4°C or at -20°C if preferred.
Storage in frost-free freezers is not recommended.
This product should be stored undiluted. Avoid repeated freezing and thawing.

Shelf Life

Please see label for expiry date.

Acknowledgements

Proclin™ 300 is a trademark of The Dow Chemical Company (“Dow”) or an affiliated company of Dow.

Health And Safety Information

Material Safety Datasheet documentation #10265 available at:
10265: <https://www.bio-rad-antibodies.com/uploads/MSDS/10265.pdf>

Regulatory

For research purposes only

Related Products

Recommended Useful Reagents

[ELISA BSA BLOCK \(BUF032A\)](#)

[ELISA ULTRABLOCK \(BUF033A\)](#)

[ELISA SYNBLOCK \(BUF034A\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M317063:180618'

Printed on 20 Jun 2018