

Datasheet: HCA053

Description:	HUMAN ANTI Ki67
Specificity:	Ki67
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	AbD02531
Isotype:	HuCAL/mouse IgG1 Fc
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (1)	▪			0.2 ug/ml - 1.0 ug/ml
ELISA	▪			2.0 ug/ml
Western Blotting	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species	Human
Species Cross Reactivity	Based on sequence similarity, is expected to react with: Bovine, Macaque, Dog, Chimpanzee, Rhesus Monkey N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Chimeric human-mouse IgG1 antibody selected from the HuCAL® phage display library and expressed in a human cell line. The antibody has variable regions of human origin and an Fc portion (including CH1 and CL domains) from mouse IgG1. It can be detected by anti mouse Fc specific secondary antibodies. This antibody is supplied liquid.
Preparation	Purified IgG prepared by affinity chromatography on Protein A
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.01% Thiomersal
Approx. Protein	IgG concentration 0.5 mg/ml

Concentrations

Immunogen Peptide derived from the human Ki67 protein, sequence GFKELFQTPG, coupled via a C-terminal cysteine to carrier proteins

External Database Links

UniProt:

[P46013](#) [Related reagents](#)

Entrez Gene:

[4288](#) MKI67 [Related reagents](#)

Specificity

Human anti Ki67 antibody, clone AbD02531 recognizes the Ki67 cell-cycle associated protein. Ki67 is expressed in proliferating cells but not in quiescent cells. Expression of this antigen occurs preferentially during late G1, S, G2, and M phases of the cell cycle, while in cells in G0 phase the antigen cannot be detected. Consequently, Ki-67 antigen expression is used in tumor pathology to detect proliferating cells in neoplastic diseases. In cultured cells, Ki-67 is expressed in the nucleolus of interphase cells. The Ki67 gene contains 15 exons. The Ki67 repeat region, within which there is a 22-amino acid Ki67 motif, is encoded by exon 13. The shorter isoform lacks exon 7. Northern blot analysis reveals multiple transcripts ranging from approximately 8.9 to 12.5 kb in proliferating but not quiescent cells. Immunoblot analysis shows expression of 320 and 359 kDa proteins. Antisense oligonucleotides inhibit cellular proliferation in a dose-dependent manner, suggesting that Ki67 protein expression may be an absolute requirement for cell proliferation.

Within cells Ki67 is predominantly localized in the G1 phase in the perinucleolar region, in the later phases it is also detected throughout the nuclear interior, being predominantly localized in the nuclear matrix. In mitosis, it is present on all chromosomes.

Histology Positive Control Tissue

Human tonsil

References

1. Jarutat, T. *et al.* (2006) Isolation and comparative characterization of Ki-67 equivalent antibodies from the HuCAL phage display library. [Biol. Chem. 387: 995-1003.](#)
2. Liu, S.K. *et al.* (2011) Delta-like ligand 4-notch blockade and tumor radiation response. [J Natl Cancer Inst. 103 \(23\): 1778-98.](#)

Storage

Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

12 months from date of despatch

Acknowledgements

Sold under license of U.S. Patents 6753136, 7785859 and 8273688 and corresponding patents. This antibody was developed by Bio-Rad, Zeppelinstr. 4, 82178 Puchheim, Germany.

Health And Safety Information

Material Safety Datasheet documentation #10094 available at: 10094: <https://www.bio-rad-antibodies.com/uploads/MSDS/10094.pdf>

Licensed Use

For *in vitro* research purposes only, unless otherwise specified in writing by Bio-Rad.

Regulatory

For research purposes only

Technical Advice

Recommended protocols and further information about HuCAL recombinant antibody technology can be found in the [HuCAL Antibodies Technical Manual](#)

Related Products

Recommended Secondary Antibodies

Goat Anti Human IgG F(ab')₂ (0500-0099...) [HRP](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M312405:180215'

Printed on 20 Jun 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)