

Datasheet: MCA4660

Description:	HAMSTER ANTI MOUSE CD137
Specificity:	CD137
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	17B5
Isotype:	IgG
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
External Database Links	<p>UniProt: P20334 Related reagents</p> <p>Entrez Gene: 21942 Tnfrsf9 Related reagents</p>
Synonyms	Cd137, Ila, Ly63

Specificity **Hamster anti Mouse CD137 antibody, clone 17B5** recognizes mouse CD137, also known as 4-1BB ligand receptor, a member of the tumour necrosis factor superfamily. CD137 is a potent costimulatory receptor for the growth and functional maturation of activated T-cells in the presence of T-cell receptor signal. It is found on activated T-cells, NK cells, monocytes and dendritic cells. Signalling through CD137 triggers T-cell expansion, cytokine production and prevention of activation-induced death of effector T-cells. CD137-mediated activation is distinct from and independent of CD28. Antibody cross-linking of CD137 on NK cells stimulates the secretion of IFN-gamma, cellular proliferation, CD25 expression and responsiveness to IL-2 and is associated with tumour activity, but not cytotoxicity.

Hamster anti Mouse CD137 antibody, clone 17B5 may be used to block CD137-mediated T-cell proliferation.

Flow Cytometry Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.

References 1. Zheng, G. *et al.* (2004) The 4-1BB costimulation augments the proliferation of CD4+CD25+ regulatory T cells. [J Immunol. 173 \(4\): 2428-34.](#)

Storage Store at +4°C or at -20°C if preferred.
Storage in frost-free freezers is not recommended.
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 18 months from date of despatch.

Acknowledgements This product may be used for research purposes only. It is not licensed for resale and may only be used by the buyer. This product may not be used and is not licensed for clinical assays, where the results of such assays are provided as a diagnostic service. If a diagnostic or therapeutic use is anticipated, then a license must be requested from the University of California. The availability of such diagnostic and therapeutic use license(s) cannot be guaranteed from the University of California.

Health And Safety Information Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Hamster IgG (STAR104...) [DyLight®549](#), [DyLight®649](#), [DyLight®800](#),
[FITC](#)

Goat Anti Hamster IgG (STAR79...) [Biotin](#), [FITC](#), [HRP](#)

Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL \(MCA2356\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

Printed on 05 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)